

Cancer and tobacco use

Tobacco use causes many cancers

Tobacco use is the leading preventable cause of cancer and cancer deaths. It can cause not only lung cancer — but also cancers of the mouth and throat, voice box, esophagus, stomach, kidney, pancreas, liver, bladder, cervix, colon and rectum, and a type of leukemia. Each year, 660,000 people in the US are diagnosed with and 343,000 people die from a cancer related to tobacco use. We have made progress: more than 1 million tobacco-related cancer deaths have been avoided since 1990 because of comprehensive cancer and tobacco control programs, early detection of cancer, and improvements in cancer treatment. However, not all states or all people have experienced the benefits of these efforts. When states make greater and longer investments in comprehensive cancer and tobacco control programs, fewer people use tobacco and get or die from tobacco-related cancers.

States and communities can

- Support comprehensive cancer control programs focusing on cancer prevention, education, screening, access to care, support for cancer survivors, and good health for all.
- Fund comprehensive tobacco prevention and control programs at CDC-recommended levels.
- Make tobacco cessation treatments more available to people who want to quit.
- Protect nonsmokers from secondhand smoke in all indoor public places and worksites, including all restaurants, bars, and casinos.

Want to learn more? www.cdc.gov/vitalsigns/cancerandtobacco

12

Tobacco use causes at least 12 types of cancer.

40%


Cancers linked to tobacco use make up 40% of all cancers diagnosed.

3 in 10

Cigarette smoking causes 3 in 10 of all cancer deaths.



Centers for Disease
Control and Prevention
National Center for Chronic
Disease Prevention and
Health Promotion



Problem:

People are still dying from cancers caused by tobacco use.




People who use tobacco or are exposed to secondhand smoke are more likely to get and die from cancer.

- Tobacco smoke has at least 70 chemicals that cause cancer, also known as carcinogens.
- Lung and colorectal cancers make up more than half of all cancers linked to tobacco use.
- Secondhand smoke exposure causes about 7,300 lung cancer deaths among nonsmoking adults each year.

Tobacco use is the leading preventable cause of cancer and cancer deaths.

- About 3 in 10 cancer deaths are caused by cigarette smoking. Lung cancer is the leading cause of cancer death for both men and women.
- Quitting tobacco use at any age can reduce the risk of getting or dying from cancer.
- Getting screened for cancer can lead to fewer people getting or dying from some tobacco-related cancers (cervix, colorectal, and lung).

Tobacco use* causes cancer throughout the body.



Tobacco-related cancers are higher among some groups:


Men



African Americans



People living in counties with low education




People living in counties with high poverty




* Tobacco use includes smoked (cigarettes and cigars) and smokeless (snuff and chewing tobacco) tobacco products that, to date, have been shown to cause cancer.

Prevent cancer deaths from tobacco use.

Tobacco-related cancer deaths have decreased over time.



Tobacco-related cancers across the United States.



Tobacco-related cancers per 100,000 persons



SOURCE: National Vital Statistics System. www.cdc.gov/nchs/nvss

More data available at: www.cdc.gov/uscs

ABOUT 1.3 MILLION tobacco-related cancer deaths have been avoided since 1990.

CDC funds 65 Comprehensive Cancer Control Programs (US states, DC, tribes, and territories) to:

PREVENTION

CREATE

tobacco-free environments to help prevent cancer.

EARLY DETECTION AND CARE

INCREASE

access to early detection and care for tobacco-related cancers.

SURVIVORSHIP

HELP

cancer survivors who use tobacco quit.

GOOD HEALTH FOR ALL

IMPROVE

cancer outcomes for everyone, especially communities with higher tobacco-related cancers and deaths.

What Can Be Done?



The Federal government is

- Requiring most insurance plans to cover recommended cancer screenings, vaccinations, as well as counseling and medications to help people quit tobacco use at no cost to the patient through the Affordable Care Act.
- Funding state tobacco and cancer prevention and control programs to help reduce the number of people who get cancer caused by tobacco use.
- Funding programs that increase cervical and colorectal cancer screening rates.
- Regulating manufacturing, marketing, and distribution of tobacco products.
- Encouraging smokefree places that protect nonsmokers from secondhand smoke.
- Funding and promoting educational campaigns about tobacco use, information, and proven resources to help people quit tobacco use.

States and communities can

- Support comprehensive cancer control programs focusing on cancer prevention, education, screening, access to care, support for cancer survivors, and good health for all.
www.cdc.gov/cancer/ncccp
- Fund comprehensive tobacco prevention and control programs at CDC-recommended levels.
www.cdc.gov/tobacco/stateandcommunity/best_practices
- Make tobacco cessation treatments more available to people who want to quit.
- Protect nonsmokers from secondhand smoke in all indoor public places and worksites, including all restaurants, bars, and casinos.

Healthcare providers can

- Ask all patients whether they use tobacco products, advise those who do not use them not to start, encourage those who do to quit, and provide help with quitting.
- Help patients understand what cancer screening tests are needed and are best for them. Make sure their patients are screened on time.
- Let patients know that there is no risk-free level of exposure to secondhand smoke and advise them to avoid it.
- Know which screening and counseling services are reimbursable to their practice.

Everyone can

- Quit using tobacco or never start. Quitting at any age will help you avoid getting cancer. Ask a healthcare provider for help quitting and call [1-800-QUIT-NOW](tel:1-800-QUIT-NOW) or go to www.smokefree.gov for free assistance.
- Encourage friends, family, and coworkers to quit using tobacco.
- Teach children and adolescents about the health risks of tobacco use and secondhand smoke exposure.
- Make their home and vehicle 100% tobacco free and not allow anyone to use any tobacco products around children.
- Learn what cancer screening tests and vaccines are right for them and their family.

1-800-CDC-INFO (232-4636)

TTY: 1-888-232-6348

www.cdc.gov

Centers for Disease Control and Prevention

1600 Clifton Road NE, Atlanta, GA 30329

Publication date: 11/10/2016